

Prevalence and molecular identification of *Cryptosporidium spp* in cattle in Baghdad province, Iraq

H.H. Alseady* and M.H. Kawan

* Department of Parasitology, College of Veterinary Medicine, University of Baghdad, Baghdad, Iraq
alseadyhaider@gmail.com

(Received November 12, 2018; Accepted January 13, 2019)

Abstract

The objective of this study was to record the prevalence rate of *Cryptosporidium* and to determine the *Cryptosporidium spp* in cattle in different parts of Baghdad province. A total number of 100 fecal samples from different age groups were collected. Conventional method using modified Ziehl-Neelsen for staining fecal smears and molecular techniques for detection the prevalence and determines the species of *Cryptosporidium* that cause infection. The overall prevalence of infection with *Cryptosporidium* was 21% (21/100) by conventional method; nested PCR was done that targeting 18S rRNA gene on the same samples in which *Cryptosporidium* DNA identified in 38 samples (38%). Four species of *Cryptosporidium* in cattle were detected for the first time in Baghdad province: *C. parvum* (6/10), *C. andersoni* (2/10), *C. bovis* (1/10) and *C. ryanae* (1/10). The determination and characterization of *Cryptosporidium spp* in cattle was very important to avoid the infection to other animals and handlers and for applying control programs.

Keywords: *Cryptosporidium*, Prevalence, Cattle, Nested PCR, Sequences analysis
Available online at <http://www.vetmedmosul.com>

نسبة الانتشار والتشخيص الجزيئي لطيفلي *Cryptosporidium spp* في الأبقار في محافظة بغداد، العراق

حيدر حسين عبيد السعيد و مي حميد كوان

فرع الطفيليات، كلية الطب البيطري، جامعة بغداد، بغداد، العراق

الخلاصة

هدفت هذه الدراسة إلى تسجيل نسب انتشار طفيلي الابواغ الخبيثة وتحديد أنواعه *Cryptosporidium species* في الأبقار وفي مناطق مختلفة من محافظة بغداد. جمعت 100 عينة براز من فئات عمرية مختلفة. تم الكشف عن الطفيلي بالطرق المجهرية التقليدية باستخدام ملون زيل نيلسون المعدلة لتلوين مسحات البراز والتقنيات الجزيئية لتسجيل نسب انتشار الطفيلي وتحديد الأنواع التي تسبب العدوى. بلغت النسبة الكلية للإصابة بطفيلي الابواغ الخبيثة 21% (21/100) باستخدام الطريقة التقليدية، وأجري تحليل سلسلة البلمرة المتداخل nPCR والذي استهدف جين 18S rRNA على نفس العينات حيث حدد الحمض النووي منقوص الأوكسجين DNA للطفيلي في 38% من العينات، تم الكشف عن أربعة أنواع لطيفلي الابواغ الخبيثة في الأبقار حيث سجلت لأول مرة في بغداد: *C. parvum* (6/10)، *C. andersoni* (2/10)، *C. bovis* (1/10) و *C. ryanae* (1/10). إن تحديد وتوصيف أنواع طفيلي الابواغ الخبيثة في الماشية مهما لتجنب الإصابة في الحيوانات الأخرى وكذلك العاملين واقتراح برامج السيطرة.

Introduction

Cryptosporidium spp. are obligate intracellular (extra-cytoplasmic) protozoan parasites that can infect a wide range of animals and human hosts worldwide (1). Cattle are

considered to be the major zoonotic reservoir for humans (2). Importantly, cryptosporidiosis in cattle causes acute or chronic gastrointestinal disturbance which is recognized as one of the major causes of profuse watery diarrhea, resulting in weight loss and growth retardation, reduced

milk production, morbidity and, in severe cases, death, leading to considerable economic losses (3). Cryptosporidiosis is transmitted mainly by the fecal-oral route, mostly by food or water contaminated with oocysts, by contact to infected animals or accidentally in laboratories (4). The conventional methods for the diagnosis of infection relies on the microscopic detection of *Cryptosporidium* oocysts in fecal samples, but these methods are unable to distinguish the different *Cryptosporidium* spp. based on morphometric or other phenotypic characteristics, due to lack of differentiating morphological features (5). Therefore, the molecular diagnostic techniques based on PCR can detect and identify genotypes. The molecular tools can identify more precise definitions of host-adapted specificity, transmission pathways, and zoonotic possibility of *Cryptosporidium* spp. (6). Worldwide, cattle are commonly infected with 4 *Cryptosporidium* spp. which include *C. parvum*, *C. andersoni*, *C. bovis* and *C. ryanae* (1,7). Given this diversity of *Cryptosporidium* species in cattle and differing zoonotic potential, it is important to be able to estimate their prevalence and distribution in particular geographic regions using molecular tools (8).

The objective of this study was to investigate the prevalence of *Cryptosporidium* infections and determination molecular characterization of *Cryptosporidium* spp. isolates from different ages of cattle in Baghdad province/Iraq.

Materials and methods

A total of one hundred fecal samples of cattle were collected from different age groups and both sexes from different parts of Baghdad province, during the period from the beginning of January 2018 to the end of September 2018, fecal smears were prepared from each sample, stained with modified Ziehl-Neelsen staining technique as primary diagnosis of *Cryptosporidium* oocysts (9).

Molecular diagnosis using Nested PCR (nPCR)

The nested PCR technique was performed for detection *Cryptosporidium* spp. based 18S ribosomal rRNA gene from cattle fecal samples. This method was carried out according to method described by (10), which included DNA extraction fecal samples by using AccuPrep® stool DNA Extraction Kit, (Bioneer, Korea). Primary PCR master mix preparation by using first primer pair: forward (5-AGACGGTAGGGTATTGGCCT -3) and reverse (5-TACGAATGCCCCCAACTGTC-3) (Maxime™ PCR PreMix Kit (*i*-Taq)), then placed in PCR Thermocycler (MyGene, Bioneer, Korea). Secondary PCR master mix was prepared by using second primer pair: forward (5-ATTGGAGGGCAAGTCTGGTG -3) and reverse (5 - TACGAATGCCCCCAACTGTC-3) (Maxime™ PCR

PreMix Kit (*i*-Taq)), then placed in PCR Thermocycler (MyGene, Bioneer, Korea).

Sequence analysis

The genetic analysis done by phylogenetic tree analysis between local *Cryptosporidium* spp. cattle isolates and NCBI-Blast submission *Cryptosporidium* species. Then the identification species isolates were submitted into of NCBI-GenBank. The DNA sequencing analysis was conducted by using Molecular Evolutionary Genetics Analysis version 6.0 (Mega 6.0) and Multiple sequence alignment analysis of the partial small subunit ribosomal rRNA gene based ClustalW alignment analysis and the evolutionary distances were computed using the Maximum Composite Likelihood method by phylogenetic tree UPGMA method (11).

Statistical analysis

Statistical analyses were computer assisted using SPSS, variables were assessed by Yat's Chi-square test (12).

Results

Total infection rate of *Cryptosporidium*

Conventional microscopic (modified Ziehl-Neelsen staining) method recorded the overall prevalence rate of *Cryptosporidium* infection in cattle was 21% (21/100). On the other hand, molecular (PCR) analysis identified *Cryptosporidium* spp. infection in 38% (38/100) (Table 1). All microscopy positive specimens were also found positive by PCR. The prevalence of *Cryptosporidium* detected by the two methods was significantly different ($P \leq 0.05$).

Table 1: Total infection rate of *Cryptosporidium* infection by conventional microscopic (modified Ziehl-Neelsen staining) method and molecular (Nested PCR) techniques in cattle

Host	Fecal samples	Conventional microscopy		Molecular (Nested PCR)	
		Total No. examined	No. of positive	%	No. of positive
Cattle	100	21	21% ^B	38	38% ^A

Variation in large horizontal letters refer to significant differences at a level of $P \leq 0.05$.

Nested PCR

Genomic DNA samples obtained from cattle fecal samples were subjected to molecular analysis by nested PCR using 18S rRNA gene specific primers in order to identify the species of *Cryptosporidium*. Nested PCR of all 100 samples employed in the study exhibited distinct band of 318 bp on agarose gel confirming the presence of *Cryptosporidium* spp (Figure 1).



Figure 1: Agarose gel electrophoresis image that show the Nested PCR product analysis of small subunit ribosomal RNA gene in *Cryptosporidium spp.* from Cattle fecal

samples. Where M: Marker 2000-100 bp, lane 1-12 showed some positive *Cryptosporidium spp.* samples at 318 bp Nested PCR product size.

Cryptosporidium species in cattle

The entire 10 samples which were positive by nested PCR amplification of 18S rRNA gene of *Cryptosporidium* were successfully sequenced. The results revealed the presence of four *Cryptosporidium spp.*, namely *C. parvum*, *C. andersoni*, *C. bovis* and *C. ryanae*, all of them have identity 100%. *C. parvum* was the most prevalent species being detected in 6/10, followed by *C. andersoni* 2/ 10, *C. bovis* and *C. ryanae* had the same infection rate 1/10 of each one (Table 2 and Figure 2).

Table 2: NCBI-BLAST Homology sequence identity between local *Cryptosporidium spp.* cattle isolates and NCBI BLAST *Cryptosporidium spp.* Isolates

Local cattle <i>Cryptosporidium spp.</i> No.	Gen-Bank accession No.	NCBI BLAST Homology sequence identity		
		NCBI BLAST <i>Cryptosporidium spp.</i>	Gen-Bank accession No.	Identity (%)
1	MH885559	<i>C. parvum</i> ^a	MH341586.1	100%
2	MH885560	<i>C. ryanae</i> ^c	MF671873.1	100%
3	MH885561	<i>C. parvum</i> ^a	MH341586.1	100%
4	MH885562	<i>C. parvum</i> ^a	MH341586.1	100%
5	MH885563	<i>C. andersoni</i> ^b	KX710086.1	100%
6	MH885564	<i>C. parvum</i> ^a	MH341586.1	100%
7	MH885565	<i>C. bovis</i> ^c	KC618591.1	100%
8	MH885566	<i>C. andersoni</i> ^b	KX710086.1	100%
9	MH885567	<i>C. parvum</i> ^a	MH341586.1	100%
10	MH885568	<i>C. parvum</i> ^a	MH341586.1	100%

Variation in small vertical letters refers to significant differences at a level of $P \leq 0.05$.

Discussion

The current study determines the prevalence and genotyping of *Cryptosporidium spp.* among cattle in different regions of Baghdad province. Fecal samples collected from cattle were screened by microscopy using modified Ziehl-Neelson staining technique and nested PCR. Although microscopy is cheaper to perform and only method to indicates active infection, higher prevalence of *Cryptosporidium* was recorded by the PCR in this study. The superior sensitivity of PCR in detecting *Cryptosporidium* infection has been shown earlier clinical trial and in patients from Northern India and South Africa (13,14).

The overall prevalence of bovine cryptosporidiosis in cattle in this study was 38%. This finding was comparable with the results of Benhouda *et al.* (15) whom recorded infection rate 40% in young calves in Algeria and the 35.5% were recorded in diarrheic calves in Sudan (16).

However, the prevalence was higher than the 14.4% infection prevalence reported from dairy farms in the Qinghai-Tibetan Plateau Area in China (7), 17.0% reported from cattle tested in Poland (17) and 18.6% in cattle in Ethiopia (18). On the other hand, the infection rate was lower than the 42.85% reported in calves in Kut city (19), 47.68% in pre-weaned dairy calves in Northeastern China (3) and 52.2% in neonatal calves in Algeria (20). The differences of infection ratios could be attributed to the differences in management systems, methods of rearing, age and breed of cattle, environmental conditions, the sampling method and sample size, as well as diagnostic techniques employed in different study localities.

The results of the current molecular study showed the presence of four major *Cryptosporidium spp.* in cattle. The most prevalent species in pre and post-weaned calves was *C. parvum*, followed by *C. andersoni*, *C. bovis* and *C. ryanae*. Our results are in agreement with the results that recorded by Silverlas (21) in Swedish cattle (22) in Indian

dairy calves, in calves in northeast China by Zhang *et al* (3), in Poland (17), in farm animals in Egypt (23), calves in Sudan (16) and in dairy cattle in northeast China (24) all of them recoded the four species of *Cryptosporidium*. On the other hand, only three species *C. andersoni*, *C. bovis* and *C. ryanae* were detected in the post-weaned calves, these results recorded in Malaysia by Muhid *et al.* (25) and in north-western China Qi *et al.* (26), while Silva *et al*; Shrestha (27,28) identified three *Cryptosporidium spp.* of

cattle, *C. parvum*, *C. bovis* and *C. andersoni*. In Mumbai region of India Hingole *et al.* (29) and in China Cai *et al.* (30) were detected three *Cryptosporidium* species also, *C. parvum*, *C. bovis* and *C. ryanae* in pre-weaned dairy calves. Yap *et al.* (11) detected only two species of *Cryptosporidium*: *C. ryanae* and *C. bovis* in cattle in Malaysia also Inpankaew *et al.* (31) detected two species of *Cryptosporidium*, *C. parvum* and *C. bovis* in dairy cows in Thailand.




Figure 2: Phylogenetic tree analysis based on the partial sequence Small subunit rRNA gene in local *Cryptosporidium spp.* cattle isolates that used for *Cryptosporidium spp.* genetic identification analysis. The evolutionary distances were computed using the Maximum Composite Likelihood method by phylogenetic tree UPGMA (MEGA 6.0 version). The local *Cryptosporidium spp.* cattle isolates of No. 1, 3, 4, 6, 9 and 10 were showed closed related to NCBI-Blast *Cryptosporidium parvum* isolate (MH341586.1), The local *Cryptosporidium spp.* cattle isolates of No. 2 was showed closed related to NCBI-Blast *Cryptosporidium ryanae* isolate (MF671873.1), The local *Cryptosporidium spp.* cattle isolates of No. 5 and No.8 were showed closed related to NCBI-Blast *Cryptosporidium andersoni* isolate (KX710086.1), and the local *Cryptosporidium spp.* cattle isolates of no. 7 was showed closed related to NCBI-Blast *Cryptosporidium bovis* isolate (KC618591.1) at total genetic change (0.01-0.04%).

The species and genotypes of *Cryptosporidium* infecting cattle are known to vary according to the host age and geographical distribution (32). Given that this study focused more on calves aged <1 month to 6 months old, our study has included cattle with a broader age range <6 months to 3 years of age sampled from different regions of Baghdad province to provide a broader picture of

Cryptosporidium infections in cattle in Iraq. *C. andersoni*, *C. bovis* and *C. ryanae* are truly less pathogenic than *C. parvum*, resulting in low grade infection and lower oocyst output, which in turn reduce the infection pressure among infected animals (33). Our results are in agreement with many studies performed in other countries reported by Taha *et al*; Helmy; Thomson; Ng (33,34-35) and thus confirm a

similar age dependent pattern of infection for Sudan for the first time. *C. andersoni* infection usually occurs in older calves and adult cattle (16).

This study described the diversity of *Cryptosporidium* in cattle of all ages in Baghdad province / Iraq. Four species of *Cryptosporidium* were detected in this study; *C. parvum*, *C. bovis*, *C. ryanae* and *C. andersoni*. *Cryptosporidium* infection due to *C. parvum* and *C. andersoni* was more prevalent in cattle in the study area.

Acknowledgment

The authors gratefully acknowledge Dr. Hassan Hachim for kindly providing the PCR kits and technique and sequence analysis used in the study. We also thank the cattle owners for cooperating and their help in collecting fecal samples in this study.

References

- Ryan U, Fayer R, Xiao L. *Cryptosporidium* spp. in humans and animals: current understanding and research needs. *Parasitol.* 2014;141:1667-1685.
- Chalmers RM, Katzer F. Looking for *Cryptosporidium*: the application of advances in detection and diagnosis. *Trends Parasitol.* 2013;29: 237-251.
- Zhang W, Wang R, Yang F, Zhang L, Cao J, Zhang X, Ling H, Liu A, Shen Y. Distribution and genetic characterizations of *Cryptosporidium* spp. in pre-weaned dairy calves in Northeastern China's Heilongjiang Province. *PLoS One.* 2013;8:e54857.
- Bouزيد M, Hunter PR, Chalmers RM, Tyler KM. *Cryptosporidium* pathogenicity and virulence. *Clin Microbiol Rev.* 2013; 26(1):115-134.
- Stark D, Al-Qassab SE, Barratt JL, Stanley K, Roberts T, Marriott D, Harkness J, Ellis JT. Evaluation of multiplex tandem real-time PCR for detection of *Cryptosporidium* spp., *Entamoeba histolytica*, and *Giardia intestinalis* in clinical stool samples. *J Clin Microbiol* 2011;49:257-262.
- Checkley W, White AC, Jaganath D, Arrowood MJ, Chalmers RM, Chen XM, Fayer R. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for *Cryptosporidium*. *Lancet Infect Dis.* 2015;15:85-94.
- Abeywardena H, Jex AR, Gasser RB. A perspective on *Cryptosporidium* and *Giardia*, with an emphasis on bovines and recent epidemiological findings. *Adv Parasitol.* 2015;88:243-301.
- Yap NJ, Koehler AV, Ebner J, Tan TK, Yvonne AL, Lim YA, Gasser RB. Molecular analysis of *Cryptosporidium* from cattle from five states of Peninsular Malaysia. *Mol Cell Probes.* 2016;30 (1):39-43.
- Beaver PC, Jung RC. *Animal Agents and Vectors of Human Disease.* 5th ed. Lea and Febiger, Philadelphia, 1985, p 249.
- Paul S, Chandra D, Tewari AK, Banerjee PS, Ray DD, Raina OK, Rao JR. Prevalence of *Cryptosporidium andersoni*: a molecular epidemiological survey among cattle in India. *Vet Parasitol.* 2009;161:31-35.
- Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol.* 2013; 30: 2725-2729.
- Petrie A, Watson P. *Statistics for veterinary and animal science.* 2nd ed. Ames: Blackwell Publishing; 2006. 24-112 p.
- Uppal B, Singh O, Chadha S, Jha AK. A Comparison of Nested PCR assay with conventional techniques for diagnosis of intestinal cryptosporidiosis in AIDS Cases from Northern India. *J Parasitol Res.* 2014;2014:706105.
- Omoruyi BE, Nwodo UU, Udem CS, Okonkwo FO. Comparative diagnostic techniques for *Cryptosporidium* infection. *Molecules.* 2014;19:2674-2683. doi: 10.3390/molecules19022674
- Benhouda D, Hakem A, Sannella AR, Benhouda A, Cacciò SM. First molecular investigation of *Cryptosporidium* spp. in young calves in Algeria. *Parasite.* 2017;24:15.
- Taha S, Elmalik K, Bangoura B, Lendner M, Mossaad E, Dausgies A. Molecular characterization of bovine *Cryptosporidium* isolated from diarrheic calves in the Sudan. *Vet Parasitol Res.* 2017;116(11):2971-2979. doi: 10.1007/s00436-017-5606-8
- Rzezutka A, Kaupke A. Occurrence and molecular identification of *Cryptosporidium* spp. isolated from cattle in Poland. *Vet Parasitol.* 2013;196:301-306.
- Manyazewal A, Francesca S, Pal M, Gezahegn M, Tesfaye M, Lucy M, Teklu W, Getachew T. Prevalence, risk factors and molecular characterization of *Cryptosporidium* infection in cattle in Addis Ababa and its environs, Ethiopia. *Vet Parasitol Reg Stud Reports.* 2018;13:79-84. doi: 10.1016/j.vprsr.2018.03.005.
- Mohammed ST, Khalaf AA, Fouad H. Detection the genotyping of *Cryptosporidium parvum* isolated from human and calves and studying in vivo effect of *Sacharomyces boulardii* on parasite. *Al-Mustansiriyah J Sci.* 2016;27(3):6-12.
- Aouatif O, Belkhir A, de Lucio A, Köster PC, Djoudi M, Dadda A, Khelef D, Kaidi R, Carmena D. *Cryptosporidium*-associated diarrhea in neonatal calves in Algeria. *Vet Parasitol Reg Stud Reports.* 2018;12:78-84.
- Silverlas C. *Cryptosporidium* Infection in Dairy Cattle. Prevalence, species distribution and associated management factors [PhD dissertation]. Swedish: Swedish University of Agricultural Sciences; 2010; p: 33-41.
- Venu R. Molecular detection and typing of *cryptosporidium* in dairy calves [PhD dissertation]. Chennai: College tamil nadu veterinary and animal sciences University Chennai; 2010. 600-007 p.
- Mahfouz ME, Mira N, Amer S. Prevalence and genotyping of *Cryptosporidium* spp. in farm animals in Egypt. *J Vet Med Sci.* 2014;76(12):1569-1575. doi: 10.1292/jvms.14-0272
- Tao W, Li Y, Yang H, Song M, Lu Y, Li W. Widespread occurrence of zoonotic *Cryptosporidium* spp. and subtypes in dairy cattle from northeast China: Public health concerns. *J Parasitol.* 2018;104(1):10-17.
- Muhid A, Robertson I, Ng J, Ryan U. Prevalence of and management factors contributing to *Cryptosporidium* spp. infection in pre-weaned and post-weaned calves in Johor, Malaysia. *Exp Parasitol.* 2011;127:534-538.
- Qi MZ, YQ, Fang YQ, Wang XT, Zhang LX, Wang RJ, Du SZ, Guo YX, Jia YQ, Yao L, Liu QD, Zhao GH. Molecular characterization of *Cryptosporidium* spp. in pre-weaned calves in Shaanxi Province, north-western China. *J Medical Microb.* 2015;64:111-116.
- Silva FM, Lopes RS, Araújo-Junior JP. Identification of *Cryptosporidium* spp. and genotypes in dairy cattle in Brazil. *Rev Bras Parasitol Vet.* 2013;22(1):1-7.
- Shrestha RD. Molecular epidemiology of waterborne zoonoses in the north island of New Zealand [PhD dissertation]. New Zealand: Institute of Veterinary, Animal and Biomedical Sciences, Massey University, Palmerston North, New Zealand; 2016; 119p.
- Hingole AC, Gudewar JG, Pednekar RP, Gatne ML. Prevalence and molecular characterization of *Cryptosporidium* spp. in cattle and buffalo calves in Mumbai region of India. *J Parasit Dis.* 2017;41(1):131-136. doi: 10.1007/s12639-016-0763-8
- Cai M, Guo Y, Pan B, Li N, Wang X, Tang C, Feng Y, Xiao L. Longitudinal monitoring of *Cryptosporidium* spp. in pre-weaned dairy calves on five farms in Shanghai, China *Vet Parasitol.* 2017;241:14-19.
- Inpankaew T, Jiyipong T, Sunanta C, Kengradomkij C, Pinyopanuwat N, Jittapalpong S. Prevalence and molecular characterization of

- bovine *Cryptosporidium* from dairy cows in Northern Thailand. *Acta Parasitologica*, 2017;62(4):772-774.
32. Ng J, Yang R, McCarthy S, Gordon C, Hijawi N, Ryan U. Molecular characterization of *Cryptosporidium* and *Giardia* in pre-weaned calves in Western Australia and New South Wales. *Vet Parasitol.* 2011;176:145-150.
33. Helmy YA. Epidemiological investigations on the public health significance of *Cryptosporidium* parasites in livestock and people in the Ismailia Canal Zone of Egypt [PhD dissertation]. Freie Universität Berlin, Germany; 2014. 77 p. doi.org/10.1016/j.vetpar.2012.12.015.
34. Thomson S. Cryptosporidiosis in farm livestock [PhD dissertation]. Glasgow: University of Glasgow; 2016; P: 112-117.
35. Ng JS. Molecular epidemiology and metabolomic characterization of *Cryptosporidium* [PhD dissertation]. Murdoch: Murdoch University; 2017; p: 94.