

Molecular detection of *Trypanosoma* species in sheep and goats in Mosul city

M.S. Mahmood and W.A. Alobaidii 

Department of Microbiology, College of Veterinary Medicine, University of Mosul, Mosul, Iraq

Article information

Article history:

Received July 13, 2021
Accepted September 11, 2021
Available online March 2, 2022

Keywords:

Trypanosoma
T. congolense
T. vivax
Sheep
Goats

Correspondence:

W.A. Alobaidii
wasenamjad@yahoo.com

Abstract

In this study, we examined blood samples of 385 sheep and goats of different ages, sexes, and sources under routine microscopic examination of the blood smear (wet, thin, thick, buffy coat layer smears) to detect *Trypanosoma*. Results show that 81 samples were positive. These samples are succumbed to the molecular detection of *Trypanosoma* and other species by the extraction of parasitic DNA this parasitic DNA is detected in samples using KIN1, KIN2, and AITSF, AITSR primers. After that, conventional polymerase chain reaction was applied, and the results showed that 81 samples had a positive reaction in using KIN1 and KIN2 primers, while the positive samples were 76 when using AITSF, AITSR primers. Moreover, results showed a high rate of infection in sheep as compared with goats using both pairs of primers and two species of *Trypanosoma* in sheep and goats. Molecular was recorded, which include *T. congolense* and *T. vivax*. Animals more than 1-2 years old group showed a high rate of infection as compared with other ages group, and females have recorded a high rate of infection as compared with males. According to the source of animals, imported animals showed a high infective rate compared to native ones. This study is the first recorded *Trypanosoma* species in small ruminants in Mosul city.

DOI: [10.33899/ijvs.2021.130488.1835](https://doi.org/10.33899/ijvs.2021.130488.1835), ©Authors, 2022, College of Veterinary Medicine, University of Mosul.
This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Trypanosomosis is a protozoal microorganism caused by unicellular flagellated protozoa. It is classified with the Protozoa Sub-kingdom, Sarcocystidophora Phylum, Kinetoplastida Order, *Trypanosomatidae* Family, and *Trypanosoma* Genus: the genus *Trypanosoma* has two main groups: Stercoraria and Salivaria, which is found in blood, different body tissues, and fluids of the vertebrates (1). There are three subgenera of *Trypanosoma* that affect health: *Vivax*, *congolense*, and *Brucei* (2). *Trypanosoma* is transmitted by flies' bites because the infective stages of the parasites are found in the mouth of the infected insect vector. *Trypanosoma*'s species, namely: *T. congolense*, *T. vivax*, and *T. brucei* affects small and large ruminants (3). The intensity of the *Trypanosoma* infection depends on the animal's *spp*, age of animals, and *Trypanosoma spp*, so the pathogenesis of Trypanosomosis differs according to the

spp causing the disease (4). The main clinical signs are fever, anemia, loss of body condition, enlargement of lymph nodes, and abortion in pregnant animals. Diagnosis of *Trypanosoma* occurs by classical methods of microscopic examination of the blood smear (5). The microscopic examination does not detect the species and does not investigate multiple infections (6). PCR has pliable the amplification of specific DNA sequences. This method can be developed to detect many types of parasites. (7). The current study has been done because no molecular studies of detection the species of *Trypanosoma* in sheep and goats in Nineveh province.

Materials and methods

Animals and samples

Three hundred and eighty-five sheep and goats with different ages, sex, and source were examined under routine

microscopic examination of the stained blood smear (wet, thin, thick and buffy coat layer smears) to detect of *Trypanosoma*. There were 81 positive samples. Then, molecular detection was done. Five ml of blood was collected from the jugular vein of each animal. The blood was transported to a tube which contained EDTA then stored in -20 °C.

DNA extraction and PCR protocol

DNA was extracted from the blood by using DNA blood extraction kit (Qiagen), the method of DNA extraction was done according to the manufacturer's

instructions manuals. The first step of PCR was done to detect the *Trypanosoma* group. KIN primers universal Trypanosome can be used according to (8). PCR was performed in 25 ul volumes. The second step can be done according to Alex *et al.* (9) using AITSF and AITSR primers (Table 1). The amplification condition of PCR was done using a thermocycler (Table1). The final PCR products were separated using 1.5% agarose gel electrophoresis. Bands observed corresponded to *T. congolense* (Kilifi/Forest and Savannah); 650-800 bp, *T. brucei*; 520-540 bp, *T. simiae*; 440-500 bp, *T. Godfrey*; 320-400 bp, and *T. vivax*; 290-400 bp (9) (Figures 1 and 2).

Table 1: Primers and amplification condition which used in this study

Primer	Sequence	Amplification condition	Ref
Kin1	5'-GCGTTCAAAGATTGGGCAAT-3'	94 °C for 3min, then 94 °C 45 seconds, 68 °C 60 seconds, 72 °C 60 seconds for 35 cycle, final extension 72°C (10 min)	8
Kin2	5'-CGCCCGAAAGTTCACC-3'		
AITSF	5'-CGGAAGTTCACCGATATTGC-3'	95°C for 10 min, 37 cycles: 95°C for 30 sec, annealing at 60°C for 1 min, 72°C for 2 min, final extension at 72°C for 10 min.	9
AITSR	5'-AGGAAGCCAAGTCATCCATC -3'		



Figure 1: Amplification of KIN1 and KIN2 primers, M=marker, 1-13 (650-700bp) positive to *Trypanosoma*, 14 Negative control.




Figure 2: Amplification of AITSF and AITSR primers, M=marker, 1,2,4,6-14 positive to *T. congolense* (700bp) 3,5 positive to *T. vivax*. (300-400bp) 15 negative control.

Results

Polymerase Chain Reaction results showed 100% of infection using Kin1 and Kin2 primers, while the percentage of infection was 93.8% using AITSF and AITSR primers in the same samples (Table 2). The PCR recorded a high infection rate in sheep and goats using both pairs of primers and AITSF, AITSR showed two species of *Trypanosoma* in sheep and goats (Table 3). According to age, the high prevalence rate of infection in sheep and goats is more than 1-2 years old, while the animals recorded a lower infection prevalence for more than two years (Table 4). Females of sheep and goats recorded high prevalence rate of infection with trypanosoma than males (Table 5). The high prevalence rate of infection with *T. congolense* in imported sheep and goats when compared with native once (Table 6).

Table 2: Showed infection with *Trypanosoma* using two pairs of primers

Primer	No	No +ve	infection %
Kin1/Kin2	81	81	100
AITSF/AITSR	81	76	93.8

Discussion

Trypanosomiasis in animals is a significant problem to livestock development. Some rural areas of Africa affected by this disease. During the last few years, several researchers have detected many species in farm animals (10). In this study, the 81 samples detected using the routine microscopic method revealed that it could be positive using Kin1 and Kin2 primers, which can detect the *Trypanosoma* group, and 76 from 81 samples gave a

positive result using AITSF and AITSR primers by the PCR technique. This primer recorded two species of *Trypanosoma* in sheep and goats with a high infective rate in sheep as compared with goats. Konnai *et al.* (11) reported that 18.3% percentage of infection in livestock of trypanosomes by using KIN primers and, he studied these primers' ability to investigate and differentiate the *Trypanosoma* using a single PCR (12). In this study, only two spp of *Trypanosoma* were recorded in sheep, including *T. congolense* and *T. vivax*, so *T. congolense* is predominant. Several studies take several spp of *Trypanosoma* in small ruminants (13) showed that *T. vivax* was the most dominant *Trypanosoma* in animals with an infection rate reached 20.91%. *T. simiae* was rarely detected, only two goats and

one sheep. At the same time, Bourzat and Gouteux (14) showed *T. vivax* with a high infection rate with clinical signs in animals, the increased prevalence of *T. vivax* approved by Authié *et al.* (15), which may result from the virulent of *Trypanosoma*, which is low and better controlled by animals, or from the mechanical transmission which has not recorded in the other spp, except *T. congolense*. The lower prevalence of *T. congolense* "forest type" when compared with *T. vivax* in domestic animals approved by Sidibe *et al.* (16); Bengaly *et al.*, (17), which is due to the increased parasitemia in *T. congolense* with anemia which leads to the death of the animal. Another study showed that *T. vivax*, *T. brucei*, and *T. congolense* were absent in sheep but present in goats with low infection rates.

Table 3: Infection rate of *Trypanosoma* species in sheep and goats

Type	No	No +ve (%)						AITSF/AITSR
		<i>Trypanosoma</i>	<i>T. congolense</i>	<i>T. vivax</i>	<i>T. brucei</i>	<i>T. simiae</i>	<i>T. godfrey</i>	
Sheep	55	55 (67.9)	43 (56.6)	9 (11.8)	0 (0)	0 (0)	0 (0)	52 (68.4)
Goats	26	26 (32.1)	21 (27.6)	3 (4)	0 (0)	0 (0)	0 (0)	24 (31.6)

Table 4: Results of infection of *Trypanosoma* species in sheep and goats according to age

Age	Animals	No +ve (%)					
		<i>Trypanosoma</i>	<i>T. congolense</i>	<i>T. vivax</i>	<i>T. brucei</i>	<i>T. simiae</i>	<i>T. godfrey</i>
Less than one year	Sheep	10	7 (70%)	2 (20%)	0 (0)	0 (0)	0 (0)
	Goats	7	4 (57%)	2 (28%)	0 (0)	0 (0)	0 (0)
< 1-2 years	Sheep	38	32 (84.2%)	6 (15.7%)	0 (0)	0 (0)	0 (0)
	Goats	16	16 (100%)	0 (%)	0 (0)	0 (0)	0 (0)
> 2 years	Sheep	7	4 (57.14%)	1 (14.2%)	0 (0)	0 (0)	0 (0)
	Goats	3	1 (33.3%)	1 (33.3%)	0 (0)	0 (0)	0 (0)

Table 5: Results of infection rate of *Trypanosoma* species in sheep and goats according to gender

Gender	Animals	No +ve (%)					
		<i>Trypanosoma</i>	<i>T. congolense</i>	<i>T. vivax</i>	<i>T. brucei</i>	<i>T. simiae</i>	<i>T. godfrey</i>
Male	Sheep	19	13 (68.4%)	3 (15.7%)	0 (0%)	0 (0%)	0 (0%)
	Goats	9	6 (66.6%)	1 (11.1%)	0 (0%)	0 (0%)	0 (0%)
Female	Sheep	36	30 (83.3%)	6 (16.6%)	0 (0%)	0 (0%)	0 (0%)
	Goats	17	15 (88.2%)	2 (11.7%)	0 (0%)	0 (0%)	0 (0%)

Table 6: Results of infection of *Trypanosoma* species in sheep and goats according to the source of animals

Source	Animals	No +ve (%)					
		<i>Trypanosoma</i>	<i>T. congolense</i>	<i>T. vivax</i>	<i>T. brucei</i>	<i>T. simiae</i>	<i>T. godfrey</i>
Native	Sheep	20	14 (70%)	3 (15%)	0 (0%)	0 (0%)	0 (0%)
	Goats	10	7 (70%)	1 (10%)	0 (0%)	0 (0%)	0 (0%)
Imported	Sheep	35	29 (82.8%)	6 (17.1%)	0 (0%)	0 (0%)	0 (0%)
	Goats	16	14 (87%)	2 (12.5%)	0 (0%)	0 (0%)	0 (0%)

Franco *et al.* (18) and Daniel *et al.* (19) showed that the *T. vivax*, *T. congolense*, and *T. brucei* cause the infection in domestic sheep and goats, and *T. vivax* is dominant. The cause of mechanical transmission or decrease a time of

development cycle in the tsetse-fly. Wayo *et al.* (20) showed that the most dominant spp of *Trypanosomes* was *T. congolense* in ruminants, and it approved the role of small ruminants as reservoirs of parasites. The reason for

the difference between the results of the two pairs of primers could be the fact that sheep and goats may infect with other spp of Trypanosome (21), which records the infection of sheep with *T. evansi*.

Gael *et al.* (10) showed 5 *Trypanosoma spp* that infected goats (*T. vivax*, *T. simiae*, *T. simiae T. savo*, *T. congolense*, and *T. brucei*) and two spp that infected sheep (*T. simiae* and *T. theileri*) with increased trypanosome infection in both. Consequently, goats are more resistant to *Trypanosoma* than sheep, as Ng'ayo *et al.* (22) suggested. Sheep have a higher rate of infection when compared to goats and pigs. Findings show they are symmetrical with other studies that record that sheep are more infected than goats naturally (23). Another study showed that sheep were highly infected with *T. congolense* Kilifi and *T. brucei*, while pigs and goats were highly infected with *T. vivax*. (22), while Bedaso *et al.* (24) recorded 2 cases of *T. vivax* 1 sheep and one goat, rare of *T. brucei*, while Kiran and Idris (25) recorded a high prevalence rate among sheep was then the prevalence of the disease among goats.

Conclusions

Trypanosoma was affected by sheep and goats. *T. congolense* is more dominant than other species of these parasites in sheep and goats in Nineveh province.

Acknowledgment

The authors wish to thank the College of Veterinary Medicine, the University of Mosul for financially supporting this work, the veterinary teaching hospital for their support.

Conflict of interest

The authors declare no conflict of interest in the manuscript.

References

- Marc D, Alan D, De L, Zhao L, Philippe H, Sathaporn J. *Trypanosoma evansi* and *surra*: A review and perspectives on transmission, epidemiology, and control, impact, and zoonotic aspects. *Biol Med Res Inter.* 2013;2013:1-20. DOI: [10.1155/2013/321237](https://doi.org/10.1155/2013/321237)
- D'avila AM, Silva RA. Animal trypanosomiasis in south America: Current status, partnership, and information technology. *Ann New York Acad Sci.* 2000;916:199-212. DOI: [10.1111/j.1749-6632.2000.tb05291.x](https://doi.org/10.1111/j.1749-6632.2000.tb05291.x)
- Desquesnes M, Biteau F, Bouyer ML, Dia L, Foil L. Development of a mathematical model for mechanical transmission of trypanosomes and other pathogens of cattle transmitted by tabanids. *Inter J Parasitol.* 2009;39(3):333-346. DOI: [10.1016/j.ijpara.2008.07.004](https://doi.org/10.1016/j.ijpara.2008.07.004)
- Gutierrez C, Desquesnes M, Touratier L, Buscher P. *Trypanosoma evansi*: Recent outbreaks in Europe. *Vet Parasitol.* 2010;174(1-2):26-29. DOI: [10.1016/j.vetpar.2010.08.012](https://doi.org/10.1016/j.vetpar.2010.08.012)
- Al-Qayim LH, Altaie LH, Ghali LS. The study of biogenic iron oxide nanoparticles effects on iron status in male rabbits infected with *T. evansi*. *Iraqi J Vet Sci.* 2021;35(Suppl III):143-147. DOI: [10.33899/ijvs.2021.132058.2039](https://doi.org/10.33899/ijvs.2021.132058.2039)
- Enock M, Claire MM, Peter W, Annah K, Alex B, Joseph M. Hemiparasitic infections in cattle from a *Trypanosoma brucei rhodesiense* sleeping sickness endemic district of eastern Uganda. *Trop Med Infect Dis.* 2020;5:24. DOI: [10.3390/tropicalmed5010024](https://doi.org/10.3390/tropicalmed5010024)
- Wicher G. The diagnosis of *Trypanosoma evansi* and its immunosuppressive effect in water buffaloes and pigs. *Lab Voor Parasit Ziekten.* 2003;7:39-40. [\[available at\]](#)
- McLaughlin GL, Ssenyonga SS, Nanteza E, Rubaire-Akiki, Wafula O, Hansen RD, Vodkin MH, Novak RJ, Gordon VR, Montenegro-James S, James M, Aviles H, Armijos R, Santrich C, Weigle K, Saravia N, Wozniak E, Gaye O, Mdachi R, Shapiro SZ, Chang KP, Kakoma I. PCR- based detection and typing of parasites. Wallingford: CAB international; 1996. 261-287 p. [\[available at\]](#)
- Alex KG, Junya YD, Axel M, Kyoko H, Naoko K, Megasari M, Chihiro S. A single test approach for accurate and sensitive detection and taxonomic characterization of Trypanosomes by comprehensive analysis of internal transcribed spacer 1 amplicons. *PLoS Negl Trop Dis.* 1996;13(2):1-20. DOI: [10.1371/journal.pntd.0006842](https://doi.org/10.1371/journal.pntd.0006842)
- Gael DM, Larson B, Emmanuella J, Ologui ME, Linda BK, Telstar GN, Brice K, Jacques FM. Frequency and diversity of trypanosomes in sheep and goats from Mongo county in south Gabon, central Africa. *Vet World.* 2020;13(11):2502-2507. DOI: [10.14202/vetworld.2020.2502-2507](https://doi.org/10.14202/vetworld.2020.2502-2507)
- Konnai S, Mekata H, Odbileg R, Simuunza M, Chembensof M, Witola W H, Tembo M E, Chitambo H, Inoue N, Onuma M, Ohashi K. Detection of *Trypanosoma brucei* in field-captured tsetse flies and identification of host species fed on by the infected flies. *Vect Born Zoon Dis.* 2008;8:1-9. DOI: [10.1089/vbz.2007.0223](https://doi.org/10.1089/vbz.2007.0223)
- Desquesnes M, Davila A. Applications of PCR-based tools for detection and identification of animal trypanosomes: A review and perspectives. *Vet Parasitol.* 2002;109(3-4):213-231. DOI: [10.1016/s0304-4017\(02\)00270-4](https://doi.org/10.1016/s0304-4017(02)00270-4)
- Nimpaye H, Njiokou F, Njine T, Njitchouang GR, Cuny G, Herder S, Asonganyi T, SIMO G. *Trypanosoma vivax*, *T. Congolense* forest type and *T. simiae*: Prevalence in domestic animals of sleeping sickness foci of Cameroon. *Parasite.* 2011;18:171-179. DOI: [10.1051/parasite/2011182171](https://doi.org/10.1051/parasite/2011182171)
- Bourzat D, Gouteux JP. Données préliminaires sur le contact glossines-petits ruminants dans le massif forestier de Mayombé, Congo. *Rev d'Elevage Med Vet des Pays Tropicaux.* 1990;43(2):199-206. DOI: [10.19182/remvt.8852](https://doi.org/10.19182/remvt.8852)
- Authié E, Bringaud F, Bakalara N, Tetaud E, Baltz T. *Trypanosomoses humaines* et animales: Maladie du sommeil et Nagana. *Ann Institut Pasteur.* 1999;10(1):27-50. DOI: [10.1016/S0924-4204\(99\)80021-3](https://doi.org/10.1016/S0924-4204(99)80021-3)
- Sidibe I, Bengaly Z, Boly H, Ganaba R, Desquesnes M, Sawadogo L. Differential pathogenicity of *Trypanosoma congolense* subgroup: Implication for the strategic control of Trypanosomiasis. *Integrat Cont of Pathol.* 2002;17(6):33-35. [\[available at\]](#)
- Bengaly Z, Sidibe I, Ganaba R, Desquesnes M, Boly H Sawadogo L. Comparative pathogenicity of three genetically distinct types of *Trypanosoma congolense* in cattle: Clinical observation and hematological changes. *Vet Parasitol.* 2002;108:1-19. DOI: [10.1016/s0304-4017\(02\)00164-4](https://doi.org/10.1016/s0304-4017(02)00164-4)
- Franco JR, Simarro PP, Diarra A, Jannin JG. Epidemiology of human African trypanosomiasis. *Clin Epidemiol.* 2014;6:257-275. DOI: [10.2147/CLEP.S39728](https://doi.org/10.2147/CLEP.S39728)
- Daniel AD, Joshua JO, Kalejaiye JO, Kalejaiye L, Dada AJ. Prevalence of trypanosomiasis in sheep and goats in a region of northern Nigeria. *Revue Elev Med Vet Pays Trop.* 1994;47(3):295-297. [\[available at\]](#)
- Wayo B, Samdi SM, Fajinmi AO, Bizi R, Dauda H, Muhammad AA, Kalejaiye JO. Prevalence of trypanosomiasis in sheep in the kachia grazing reserve. *African J Clin Experiment Microbiol.* 2017;18:159-170. DOI: [10.4314/ajcem.v18i2.10](https://doi.org/10.4314/ajcem.v18i2.10)
- Suheir E, Abdelmajeed NA, Al-Jawabreh T, Hanan A, Nahed, A, Ziad A. Prevalence of *Trypanosoma evansi* in livestock in Palestine. *Parasit Vect.* 2020;13(21):1-8. DOI: [10.1186/s13071-020-3894-9](https://doi.org/10.1186/s13071-020-3894-9)
- Ngayo MO, Njiru ZK, Kenya EU, Muluvi GM, Osir EO, Masiga DK. Detection of trypanosomes in small ruminants and pigs in Western

- Kenya: Important reservoirs in the epidemiology of sleeping sickness. *Kinetoplastid Biol Dis.* 2005;4(1):5. DOI: [10.1186/1475-9292-4-5](https://doi.org/10.1186/1475-9292-4-5)
23. Masiga DK, Okech G, Irungu P, Ouma J, Wekesa S, Ouma B, Guya SO, Ndungu JM. Growth and mortality in sheep and goats under high tsetse challenge in Kenya. *Trop Anim Health Pro.* 2002;34:489-501. DOI: [10.1023/a:1021241220575](https://doi.org/10.1023/a:1021241220575)
24. Bedaso K, Seifu H, Getachew T. Prevalence and pathogenic significance of trypanosomosis on sheep and goats of Mareka district, Dawro zone, Southwestern Ethiopia. *J Anim Sci Livestock Prod.* 2017;1:102. DOI: [10.21767/2577-0594.100002](https://doi.org/10.21767/2577-0594.100002)
25. Kiran S, Abdurrahman I. Prevalence of trypanosomiasis among sheep and goats slaughtered at Sokoto central abattoir. *Inter J Anim Sci Husb Livestock Product.* 2017;3(11):233-236. [\[available at\]](#)

التقصي الجزيئي عن أنواع المثقبيات في الضأن والماعز في مدينة الموصل

مروة سمير محمود و وسن امجد العبيدي

فرع الأحياء المجهرية، كلية الطب البيطري، جامعة الموصل، الموصل، العراق

الخلاصة

تم من خلال هذه الدراسة فحص عينات دم ٣٨٥ رأساً من الضأن والماعز من مختلف الأعمار والأجناس والمصدر والتي تم فحصها تحت الفحص المجهرى الروتيني من خلال فحص المسحات الدموية (الرطبة، الخفيفة، السمكية، مسحة طبقة الخلايا اللمفية) وذلك للكشف عن طفيلي المثقبيات والتي أظهرت أن ٨١ عينة كانت موجبة، تم إخضاع هذه العينات الموجبة للكشف الجزيئي عن طفيلي المثقبيات والأنواع عن طريق استخلاص الحمض النووي الطفيلي، ثم تم الكشف عن هذا الحمض النووي في العينات باستخدام بادئات KIN1 و KIN2 و AITSF و AITSR ثم تم تطبيق تفاعل السلسلة المتبلورة التقليدي، وأظهرت النتائج أن ٨١ عينة أعطت تفاعلاً إيجابياً باستخدام بادئات KIN1 و KIN2 بينما عند استخدام البادئات AITSF، AITSR كان عدد العينات الموجبة ٧٦. أظهرت النتائج إصابة عالية في الضأن عند مقارنتها بالماعز باستخدام زوج من البادئات فضلاً عن أنه تم الكشف الجزيئي عن نوعين من المثقبيات في الأغنام والماعز *T. congolense* و *T. vivax*. أظهرت الحيوانات التي يزيد عمرها عن ١-٢ سنوات نسبة مرتفعة للإصابة عند مقارنتها مع الفئات العمرية الأخرى، وسجلت الإناث نسبة عالية للإصابة عند مقارنتها بالذكور. وفقاً لمصدر الحيوانات، أظهرت الحيوانات المستوردة معدل إصابة مرتفعاً عند مقارنتها بالحيوانات المحلية. وتعد هذه الدراسة هي الأولى من نوعها للكشف عن أنواع المثقبيات في المجترات الصغيرة في مدينة الموصل.